

KYLE R. WANZEL MD, MEd, FRCS(C) Plastic, Reconstructive and Cosmetic Surgeon



1243 Islington Ave., Suite 600, Toronto, ON M8X 1Y9 Phone: (416) 236-2571 Fax: (416) 236-4591 www.plasticcosmeticsurgery.ca

## **Post-Operative Instructions for Breast Lift**

- 1. Take your medication(s) as prescribed
- 2. Expect a small to moderate amount of pain, tenderness, swelling, numbness and bruising
  - i. It is not unusual for one breast to be more swollen and/or bruised than the other one
- 3. Smoking will delay the healing process
  - i. If you smoke, you must quit at least 6 weeks before and 6 weeks after your surgery
- 4. During the first week, the majority of your time should be spent relaxing, although some leisurely walking is permitted
  - i. No strenuous activity or heavy lifting with the arms is permitted for 6 weeks
- 5. Wear your support bra for 24 hours a day for the first 4 weeks post-op (this may be removed for showering)
- 6. Return in approximately 5 to 7 days for reassessment and dressing change
- 7. You may shower daily (starting after the first dressing change) and gently pat towel dry
- 8. If required, gauze, ABD or "maxi pads" may be used as fresh dressings inside your bra
- 9. Leave all paper tapes on over the incisions until the first post-op visit (you may shower with these on and get them wet). They will start to lift off on their own.
- 10. A small amount of yellow discharge from under the breasts is common early postop
  - i. It is also not unusual for there to be a very small open area under the breast in the center of the breast (this will heal in on its own without a problem)
- 11. Light activities and more walking are permitted at 1 week postop
- 12. Return again in approximately 1 week for a follow-up visit
- 13. Normal and more strenuous activities are permitted at around 6 weeks post-op

## IF YOU HAVE ANY CONCERNS AT ALL, PLEASE CONTACT THE OFFICE